

ORIGINAL ARTICLE

Correlation of Smoking Habits, Physical Activities and Fat Intake with Cognitive Ability in Indonesian Elderly

Yuniars Renowening^{1*}, Suradi Suradi², Ari Probandari²

1. Postgraduate Student of Nutrition Science, Sebelas Maret University, Ir Sutami Street No 36A Ketingan, Jebres, Surakarta 57126, Indonesia

2. Departement of Public Health, Faculty of Medicine, Sebelas Maret University, Ir Sutami Street No 36A Ketingan, Jebres, Surakarta 57126, Indonesia

ARTICLE INFO

Keywords:

Smoking
Physical activity
Cognitive ability
Fat
Nutrition

*Corresponding author:

Yuniars Renowening
Postgraduate Student of Nutrition
Science, Sebelas Maret University,
Ir Sutami Street No 36A Ketingan,
Jebres, Surakarta 57126, Indonesia.
Tel: +6285728502393

Email:

yuniarsrenowening@gmail.com

Received: April 18, 2019

Revised: September 21, 2019

Accepted: September 29, 2019

ABSTRACT

Background: The aging process that occurs in elderly makes the elderly more vulnerable to health threats, such as decreased cognitive abilities. Lifestyle is a factor that plays an important role in several diseases, such as smoking, lack of physical activity, and consuming unhealthy foods. The purpose of this study was to find the correlation of smoking habit, physical activities and fat intake with cognitive abilities in elderly.

Methods: This study used a cross sectional design involving 182 elderly over the age of 60 years in six integrated services post (Posyandu) in Klaten Regency, Indonesia. Data of smoking habits and physical activities were obtained by interview and fat intake was determined by food recall. Data of cognitive ability was obtained through interview using Mini Mental State Exam (MMSE) questionnaire. Data were tested using Spearman correlation to determine the relationship between variables.

Results: The results showed that there was a significant relationship between smoking habit with cognitive ability in elderly ($P < 0.001$). There was a significant relationship between physical activity and cognitive ability in elderly ($P < 0.001$). There was no significant relationship between fat intake and cognitive ability in elderly ($P = 0.494$).

Conclusion: Low smoking habits and high physical activity was correlated to a better cognitive function in Indonesian elderly.

Please cite this article as: Renowening Y, Suradi S, Probandari A. Correlation of Smoking Habits, Physical Activities and Fat Intake with Cognitive Ability in Indonesian Elderly. Int J Nutr Sci. 2019;4(4):186-191.

Introduction

Elderly is the last period in the cycle of human life that will be experienced later by everyone. As one gets older, physiological functions will decrease due to degenerative processes (aging), so the problems related to older age will require more attention in future (1). The aging process in elderly makes the elderly more vulnerable to various health threats (2). Elderly will experience a decrease in tissue capacity too (3) and body function will also experience setbacks, a decreased memory, and a

cognitive health decline (4, 5). Lifestyle is a factor that plays an important role in the occurrence of several diseases, such as smoking, lack of physical activities, and consuming unhealthy foods (6).

Smoking is a habit that has a huge impact on health (7). The prevalence of elderly smoking in Indonesia is relatively high. It was shown that as many as 37.5% of the elderly are active smokers who smoke an average of 13 cigarettes per day (8). Central Java is one of the provinces with the highest number of smokers. In 2017, the number of smokers reached

62.7% with an average number of 10 cigarettes per day and in Klaten Regency, the number of active smokers was 36.2% (9). Smoking is thought to have a negative effect on cognitive ability. Studies conducted in Australia, showed that seniors who smoked experienced greater cognitive decline each year than seniors who did not smoke (10). Cataldo et al.'s study (2010) revealed that the duration of smoking increased the risk factors for cognitive disability associated with increased oxidative stress and inflammation in smokers (11).

Physical activity is an indicator that can be utilized to evaluate overall health status of elderly (12, 13). Regular physical activity is beneficial to health, extends independent and active life expectancy, and improves quality of life in elderly. Aging is very influential in physical activity in elderly, for instance, the reduced ability to perform daily activities, independently (14). Scientific evidences also demonstrated that elderly people aged 60 years or older and with low physical activity had a high risk of experiencing cognitive disability, depression, and dementia (15).

Two thirds or more of diseases in elderly are closely related to nutrition. According to gerontologists and geriatrics, it is estimated that 30-50% of nutritional factors are important to achieve and maintain the elderly health (4). Fat is a nutrient that influences cognitive ability. A study has shown that there is a decrease in memory status in rats fed with high saturated fat for three months (16). Another study conducted on mice also showed that a high-fat diet lowered haptoglobin, a molecule that plays a role in the pathogenesis of neurodegenerative diseases (17).

Physical activity in the elderly over 60 years tends to be lower than the elderly with a younger age (18). In addition, the elderly also tend to have

high fat intake due to poor diet (19). High smoking habits, lack of physical activity, and high fat intake in the elderly underlie the importance of conducting a research on the correlation of smoking habits, physical activity and fat intake with cognitive ability in Indonesian elderly.

Materials and Methods

This study was a cross sectional research design and the population of study were elderly over 60 years old in Klaten Regency, Central Java (n=182). The inclusion criteria were elderly men and women over 60 years old, able to chew and live with family. The exclusion criteria were elderly with overnutrition status [Body mass index (BMI) ≥ 23.5] and to be illiterate. This study was approved by the Health Research Ethics Commission Sebelas Maret University No.74/UN27.06/KEPK/2019.

Data on smoking habits and physical activity were obtained from respondents by interview, and fat intake results were obtained using a 24-hour recall questionnaire. Classification of smokers was based on World Health Organization (WHO) report in 2013 (20), classification of physical activity was based on Physical Activity Scale for Elderly (PASE) (21), and fat intake was classified based on the 2013 Nutrient Adequacy Ratio (NAR) according to the Indonesian Ministry of Health (22). Data on cognitive ability were obtained by interview using Mini Mental State Exam (MMSE) questionnaire together with categories based on the 2010 Shigemon et al.'s study (23). The classification of these variables was presented in Table 1. SPSS software (Version 21, Chicago, IL, USA) was used for statistical analysis and the data were compared using the statistical test of bivariate analysis, and Spearman. A p value less than 0.05 was considered significant.

Table 1: Classification of the study variables

Variable	Category	Classification
Smoking habits*	<1 stick per day	Not a smoker
	1-10 sticks per day	Light smoker
	11-20 sticks per day	Moderate smoker
	>20 sticks per day	Heavy smoker
Physical activities**	>16	High
	9-16	Moderate
	<9	Low
Fat intake***	<70%	Low
	70-79%	Relatively low
	80-99%	Sufficient
	>100%	High
Cognitive ability****	<16	Low
	17-23	Moderate
	>24	Normal

*World Health Organization (WHO, 2013), ** Physical Activity Scale for Elderly (PASE), *** Nutrient Adequacy Ratio (NAR, 2013), ****Shigemon et al. (2010)

Results

The number of elderly women involved in the study was higher than men, which was 52.1%. Based on the age category, the majority of elderly were ranging from 60 to 70 years (69.2%), 71-80 years as much as 25.9% and the least number was among those older than 80 years which only reached 4.9%. Based on bivariate analysis, there was a significant relationship between smoking habits and cognitive ability ($P < 0.001$, Table 2). The correlation coefficient also showed a positive relationship between smoking and cognitive ability. Based on bivariate analysis, there was a significant relationship between physical activity and cognitive ability ($P < 0.001$, Table 3). The correlation coefficient also showed a positive relationship between physical activity and cognitive ability. Based on bivariate analysis, there was no significant relationship between fat intake and cognitive ability ($P = 0.494$, Table 4).

Discussion

In this study, there was a significant relationship of smoking habits and physical activity with cognitive abilities in the elderly, while fat intake was not significantly related with cognitive abilities in the elderly. The difference between this study and other studies was the variable of fat intake. There were no studies to discuss specifically the relationship of fat intake with cognitive abilities in elderly. Aging is

often associated with an increase in the number of chronic disease conditions and physical decline that can negatively impact health status. Lifestyle has become an important component affecting health and well-being in elderly. Most health problems can be prevented through healthy living habit such as avoiding smoking, doing physical activity and dietary modification (24).

Elderly with a nutrition-rich dietary and conducting regular physical activities was illustrated to have a lower risk of diseases. The risk of anxiety and depression can also be prevented through healthy eating and active lifestyle (25). Smoking can cause an accumulation of oxidants and free radical. Consequently, excess oxidants and free radical trigger oxidative stress. Research conducted on adults who smoke shows increased oxidative stress contributes to the formation of neurofibrillary plaques, which triggers cognitive decline (26).

In line with these findings, Wang et al. stated that oxidants and free radicals can oxidize DNA, RNA, protein, and fat which will cause damage to cells too (27). Smoking is also a risk factor for cardiovascular and cerebrovascular disease, stroke, increased oxidative stress, atherosclerosis, and inflammation which can negatively impact cognitive function and the occurrence of dementia (28). Physical activity is very useful for elderly. Regular physical activity can prevent cognitive decline and improve independent

Table 2: Correlation of smoking habit and cognitive ability in elderly

Smoking habit	Cognitive ability (182)			Correlation coefficient	P value*
	Low	Moderate	Normal		
Heavy smoker	2	3	3	0.348	<0.001
Moderate smoker	28	13	15		
Light smoker	8	14	19		
Not a smoker	3	32	42		

*Significant at $\alpha = 0.05$, using the Spearman Correlation test

Table 3: Correlation of physical activities and cognitive ability in elderly

Physical activity	Cognitive ability (182)			Correlation coefficient	P value*
	Low	Moderate	Normal		
Low	16	24	18	0.248	<0.001
Moderate	21	24	30		
High	4	14	31		

*Significant at $\alpha = 0.05$, using the Spearman correlation test

Table 4: Correlation of fat intake and cognitive ability in elderly

Fat intake	Cognitive ability (182)			Correlation coefficient	P value*
	Low	Moderate	Normal		
Low	4	9	8	0.051	0.494
Relatively low	1	1	4		
Sufficient	7	7	6		
High	29	45	61		

*Significant at $\alpha = 0.05$, using the Spearman Correlation test

life in elderly. The results of this study are in line with research conducted in Taipei on 275 elderly people showing that physical activity can increase happiness and mental health, because it can stimulate anti-anxiety effects and also improve sleep quality (29).

Physical activity is a lifestyle that can be modified and brings advantages in maintaining cognitive function as one gets older (30, 31). Physical activity is able to regenerate old cells and increase the number of young cells, thus it slows down cognitive decline (32, 33). The nutrition needs of elderly are influenced by several factors such as age, sex, physical activities, occupation, body posture and health conditions (34). The intake needed by elderly must be in accordance with the Nutrient Adequacy Ratio (NAR) to achieve a balanced nutritional status (35).

Excessive fat consumption is one of the causes of accumulation of oxidative stress and may bring damage to antioxidant defenses. In this study, there was no significant relationship between fat intake and cognitive ability in the elderly, which is in contrast to previous studies done on mice with a significant increase in lipid peroxidation along with a decrease in antioxidant activity, and this can trigger cognitive decline in the elderly (36). The Wyss-Coray et al.'s study also stated that excess fat will accumulate and cause inflammation in brain and triggers cognitive decline (37). The difference in results can be caused by bias during the recall process, a less in-depth approach to the food recall process (38).

Conclusion

The findings in this study highlighted that elderly with intense smoking habits have low cognitive ability. It was shown that elderly with high physical activities have good cognitive ability. This study showed no significant relationship between fat intake and cognitive ability in elderly. It is expected that in the future the elderly can maintain a good lifestyle by avoiding smoking, excessive fat consumption and increase in physical activity to maintain good cognitive abilities too.

Acknowledgment

The authors would like to thank the parents and friends who have helped this research. The authors also thank all the participants involved in this study and Sebelas Maret University for financial support of the research. We are also grateful to the puskesmas and enumerators who assisted this research.

Conflict of Interest

None declared.

References

- 1 Kemenkes RI. Situasi Lanjut Usia (Lansia) di Indonesia. Pusat Data dan Informasi Kementerian dan Kesehatan RI. *Kementerian Kesehatan RI*. 2016.
- 2 Lepeule J, Bind MC, Baccarelli AA, et al. Epigenetic Influences on Associations between Air Pollutants and Lung Function in Elderly Men: The Normative Aging Study. *Environmental Health Perspectives*. 2014;122:566-72. DOI:10.1289/ehp.1206458. PMID:24602767.
- 3 Abdollahzade SM, Eftekhari MH, Almasi-Hashiani A. The prevalence of malnutrition in elderly members of Jahandidegan Council, Shiraz, Iran. *Int J Nutr Sci*. 2018;3:86-91.
- 4 Ariati NN, Gumala NMYG, Nursanyoto H. Hubungan Konsumsi Makronutrien dengan Resiko Penuaan Dini pada Lansia Yang Mengikuti Senam Lansia di Posyandu Kabupaten Ganyar. *J Sangkareang Mataram*. 2017;3:34-7.
- 5 Pritasari, Damayanti D, Lestari NT. Gizi Dalam Daur Kehidupan. Badan Pengembangan dan Pemberdayaan Sumber Daya Manusia. *Kementerian Kesehatan RI*. 2017.
- 6 Mehrabani G, Aminian S, Mehrabani G, et al. Dietetic plans within the multiple sclerosis community: A review. *Int J Nutr Sci*. 2019;4:14-22.
- 7 Akbarzadeh M, Almasi-Hashiani A, Farahmand M. The prevalence of risk factors of non-communicable diseases in Fars Province. *Int J Nutr Sci*. 2016;1:23-9.
- 8 Kemenkes RI. Gambaran Kesehatan Lanjut Usia di Indonesia. Pusat Data dan Informasi Kementerian dan Kesehatan RI. Kementerian Kesehatan RI Dinas Kesehatan Provinsi Jawa Tengah. 2015.
- 9 Dinas Kesehatan Kabupaten Klaten. Profil Kesehatan Kabupaten Klaten. Dinas Kesehatan Kabupaten Klaten. *Klaten*. 2017.
- 10 Anstey KJ, Sanden C, Salim A, et al. Smoking as a Risk Factor for Dementia and Cognitive Decline: A Meta-Analysis of Prospective Studies. *Am J Epidemiol*. 2007;166:367-78. DOI:10.1093/aje/kwm116. PMID:17573335.
- 11 Cataldo JK, Prochaska JJ, Glantz SA. Cigarette Smoking is a Risk Factor for Alzheimer's Disease: An Analysis Controlling for Tobacco Industry Affiliation. *J Alzheimers Dis*. 2010;19:465-80. DOI:10.3233/JAD-2010-1240. PMID:20110594.
- 12 WHO (World Health Organisation.) Non-communicable Diseases and Mental Health. <http://www.who.int/nmh/publications/9789241597418/en/>. Accessed 2013.
- 13 Hamer M, Lavoie KL, Bacon SL. Taking up

- physical activity in later life and healthy ageing: the English Longitudinal Study of Ageing. *Br J Sports Med.* 2013;48:239-43. DOI:10.1136/bjsports-2013-092993. PMID:24276781.
- 14 Chou CH, Hwang CL, Wu YT. Effect of exercise on physical function, daily living activities, and quality of life in the frail older adults: A meta-analysis. *Arch Phys Med Rehabil.* 2012;93:237-44. DOI:10.1016/j.apmr.2011.08.042. PMID:22289232.
 - 15 Sun F, Norman IJ, While AE. Physical Activity in Older People: A Systematic Review. *BMC Public Health.* 2013;13:449. DOI:10.1186/1471-2458-13-449. PMID:23648225.
 - 16 Freeman LR, Haley-Zitlin V, Rosenberger DS, et al. Damaging Effects of a High-Fat Diet to the Brain and Cognition: A Review of Proposed Mechanisms. *Nutr Neurosci.* 2014;17:241-51. DOI:10.1179/1476830513Y.0000000092. PMID:24192577.
 - 17 Spagnuolo MS, Mollica MP, Maresca B, et al. High Fat Diet and Inflammation: Modulation of Haptoglobin Level in Rat Brain. *Front Cell Neurosci.* 2015;9:479. DOI:10.3389/fncel.2015.00479. PMID:26696835.
 - 18 Buchman AS, Wilson RS, Yu L, et al. Total Daily Activity Declines More Rapidly with Increasing Age in Older Adults. *Arch Gerontol Geriatr.* 2014;58:74-9. DOI:10.1016/j.archger.2013.08.001. PMID:24007938.
 - 19 Rohmawati N, Asdie AH, Susetyowati. Anxiety Level, Dietary Intake, and Nutritional Status of Elderly in Yogyakarta Municipality. *Jurnal Gizi Klinik Indonesia.* 2015;12:62-71.
 - 20 World Health Organization (WHO). Report on the Global Tobacco Epidemic. 2013.
 - 21 Williams K, Frei AA, Vetsch F et al. Patient-Reported Physical Activity Questionnaires: A Systematic Review of Content and Format. *Health Qual Life Outcomes.* 2012;10:28. DOI: 10.1186/1477-7525-10-28.
 - 22 Angka Kecukupan Gizi (AKG). Angka Kecukupan Gizi Energi, Protein Yang Dianjurkan Bagi Bangsa Indonesia. Lampiran Peraturan Menteri Kesehatan Republik Indonesia Nomor Tahun. 2013;2013:75.
 - 23 Shigemon K, Ohgi S, Okuyama E, et al. The Factorial Structure of The Mini Mental State Examination (MMSE) in Japanese Dementia. *BMC Geriatr.* 2010;10:36. DOI:10.1186/1471-2318-10-36. PMID:20534132.
 - 24 Mehrabani D, Vahedi M, Eftekhari MH, et al. Food avoidance in patients with ulcerative colitis: A review. *Int J Nutr Sci.* 2017;2:189-195.
 - 25 CDC (Centers for Disease Control and Prevention). The State of Aging and Health in America 2013. US Dept of Health and Human Services. Atlanta. Retrieved from <https://www.cdc.gov/aging/pdf/state-aging-health-in-america-2013.pdf>. Accessed 2013.
 - 26 Ambrose JA, Barua RS. The Pathophysiology of Cigarette Smoking and Cardiovascular Disease: An Update. *J Am Coll Cardiol.* 2004;43:1731-7. DOI:10.1016/j.jacc.2003.12.047. PMID:15145091.
 - 27 Wang CC, Lu TH, Liao WC, et al. Cigarette Smoking and Cognitive Impairment: A 10-year Cohort Study in Taiwan. *Gerontology and Geriatrics.* 2010;51:143-8.
 - 28 Aghasadeghi K, Zarei-Nezhad M, Keshavarzi A, et al. The prevalence of coronary risk factors in Iranian lor migrating tribe. *Arch Iran Med.* 2008;11:322-5. DOI:08113/AIM.0015. PMID:18426325.
 - 29 Chien MY, Kuo HK, Wu YT. Sarcopenia, cardiopulmonary fitness, and physical disability in community-dwelling elderly people. *Phys Ther.* 2010;90:1277-87.
 - 30 Yaffe K, Fiocco AJ, Lindquist K, et al. Predictors of Maintaining Cognitive Function in Older Adults the Health ABC Study. *Neurology.* 2009;72:2029-35. DOI:10.1212/wnl.0b013e3181a92c36. PMID:19506226.
 - 31 Buchman AS, Boyle PA, Yu L, et al. Total Daily Physical Activity and the Risk of AD and Cognitive Decline in Older Adults. *Neurology.* 2012;78:1323-9. DOI: 10.1212/wnl.0b013e3182535d35. PMID:22517108.
 - 32 Kim S, Ko I, Kim B, et al. Treadmill Exercise Prevents Aging Induced Failure of Memory Through an Increase in Neurogenesis and Suppression of Apoptosis in Rat Hippocampus. *Exp Gerontol.* 2010;45:357-65. DOI:10.1016/j.exger.2010.02.005. PMID:20156544.
 - 33 DeCarolis NA, Eisch AJ. Hippocampal Neurogenesis as a Target for the Treatment of Mental Illness: A Critical Evaluation. *Neuropharmacology.* 2010;58:884-93. DOI:10.1016/j.neuropharm.2009.12.013. PMID:20060007.
 - 34 Kemenkes RI. Situasi Lanjut Usia (Lansia) di Indonesia. Pusat Data dan Informasi Kementerian dan Kesehatan RI. *Kementerian Kesehatan RI.* 2016.
 - 35 Istiany A, Rusilanti. Gizi Terapan. PT Remaja Rosdakarya. *Bandung.* 2013.
 - 36 Aqababa H, Mottahedzadeh R, Shahabi S, et al. The action of ginkgo biloba on passive avoidance learning in experimental rats. *Middle East J Sci Res.* 2013;16:751-5. DOI:10.5829/idosi.mejsr.2013.16.06.7460.

-
- 37 Wyss-Coray T, Rogers J. Inflammation in Alzheimer disease: A Brief Review of the Basic Science and Clinical Literature. *Cold Spring Harb Perspect Med*. 2012;2. DOI:10.1101/cshperspect.a006346. PMID:22315714.
- 38 Althubaiti A. Information Bias in Health Research: Definition, Pitfalls, and Adjustment Methods. *J Multidiscip Healthc*. 2016;9:211-7. DOI:10.2147/JMDH.S104807 PMID:27217764.