

ORIGINAL ARTICLE

The Effect of Aqueous Extract of *Cariandrum Sativum* on Fetal Weight and Height in Pregnant Mice

Davood Mehrabani^{1,2,3}, Mahnaz Mehrmand⁴, Akbar Vahdati⁴, Mehdi Pasalar^{5*}, Mina Rabiee⁶, Seyed Jalil Masoumi^{7,8}

1. Stem Cell Technology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran
2. Burn and Wound Healing Research Center, Shiraz University of Medical Sciences, Shiraz, Iran
3. Comparative and Experimental Medicine Center, Shiraz University of Medical Sciences, Shiraz, Iran
4. Science and Research Branch, Islamic Azad University, Fars, Iran
5. Traditional Medicine Research Center, Shiraz University of Medical Sciences, Shiraz, Iran
6. Shahre-Kord Branch, Islamic Azad University, Shahre-Kord, Iran
7. Nutrition Research Center, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran
8. Department of Clinical Nutrition, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran

ARTICLE INFO

Keywords:

Cariandrum sativum
Fetus
Weight
Height
Pregnancy

***Corresponding author:**

Mehdi Pasalar,
Traditional Medicine Research
Center,
Shiraz University of Medical
Sciences,
Shiraz, Iran.
Tel: +98-71-32341025
Email: mehdipasalar@yahoo.com
Received: March 5, 2018
Revised: April 12, 2019
Accepted: April 18, 2019

ABSTRACT

Background: Herbal medicine may have irreparable effects on fetal development. This study investigated the effect of *Cariandrum sativum* (coriander) aqueous extract on mice fetal height and weight.

Methods: In an experimental study, 120 pregnant mice were divided into four equal groups. The control group was without any intervention. The experimental group received 3500, 5000 and 6500 mg/kg of aqueous extract of coriander. The height and weight of the embryos were recorded in first, second and third week of pregnancy.

Results: The height and weight of embryos in experimental groups in first, second and third week was significantly lower than the control group.

Conclusion: Coriander (*C. Sativum*) aqueous extract was shown to reduce the fetal height and weight in a dose-dependent manner. These findings can be added to the literature when coriander is going to be used during pregnancy.

Please cite this article as: Mehrabani D, Mehrmand M, Vahdati A, Pasalar M, Rabiee M, Masoumi SJ. The Effect of Aqueous Extract of *Cariandrum Sativum* on Fetal Weight and Height in Pregnant Mice. Int J Nutr Sci. 2019;4(3):137-141.

Introduction

Birth defects and congenital anomalies are similar terminologies that describe constructional, functional and metabolic disorders at birth time. Congenital defects can be caused by inherited and environmental factors, while one of the most important environmental factors is the use of herbal medicines during pregnancy (1). It was shown that pregnant women search for herbs as a natural and safe drug during pregnancy (2). In

addition to several therapeutic effects of herbs (3, 4), conventional medicine has side effects such as teratogenic and toxic impacts during pregnancy for both the mother and the fetus causing fetal abnormalities (5).

Severe uterine contractions and abortions were reported following use of herbs leading to expansion of the uterus muscles, opening of the cervix are also other side effects during pregnancy (6). For example, barberry extract (1) and *Stachys lavandulifolia vahl*

(7) were demonstrated to have decreasing effect on fetal height and weight. Ginseng was found to induce dehydration through an increase in sex hormones (5). Echium extract was shown to have a negative effect on reproductive factors and fetal development (8). The teratogenic effect of fennel extract to cause exohepaty in the fetus was previously demonstrated (9).

Coriander know as *Coriandrum Sativum L.* too belongs to *Apiaceae* family (10). Coriander's seeds contain mannitol and flavonoid glycosides such as quercetin, isoquercitrin, and rutin. There is also tannin, coumarin, mucilage, starch and chlorogenic and caffeic acid in its seeds (11). Flavonoids in plants produce blue and red pigments in flowers and protect against the attack of germs and insects. In addition, they have antioxidant, anti-allergic, anti-inflammatory, antimicrobial and anticancer activity (12).

Quercetin is known as the most active flavonoid in studies. Quercetin with its anti-inflammatory properties directly inhibits the onset of inflammatory processes. For example, it prevents the synthesis and release of histamine, a mediator of inflammatory reactions (13). Additionally, it has antioxidant activity and reduced vitamin C activity (12). Quercetin antitumor activity has also been proven (13). Recent studies have shown that quercetin can help men in the treatment of chronic prostate inflammation due to its inhibitory effects on mast cells (14).

Tannins are highly complex compounds that produce polyhydric phenols such as pyrogallol, catechole and Ellagic acid during hydrolysis (15) that can reduce irritations and pain, eliminate oral swelling, local bleeding and promote healing in bronchitis and burns (16). Coriander is used in traditional medicine for food digestion, anti-inflammation, appetite enhancer, and muscle ailments.(10) Regarding the importance of genetic and maternal abnormalities in infants and the excessive use of certain herbs during pregnancy without any knowledge on the side effects, the aim of this study was to investigate the effect of coriander extract on mice fetal height and weight.

Materials and Methods

In an experimental study, for preparation of coriander extract, after harvesting the samples and their scientific and systemic identification, the green and fresh leaves of coriander were provided. They were dried at room temperature under shadow and kept in refrigerator in glass containers until use. For preparation of aqueous extract, 100 g of the dried powder was weighed by a digital scale, and 60 mL of sterile distilled water was added while the glass container was heated to 80°C. It was placed in the

water bath at 60°C for 24 hours. Finally, the mixture was passed through a funnel and filter paper.

Determination of lethal dose of coriander was done by LD50 method. Three different doses were selected based on previous studies. Dose with 50% mortality was considered as LD50 dose defined as 6500 mg/kg. A standard two milliliters syringe equipped with a feeding needle or gavage was used for administration purposes. The skin of the neck region was stretched and kept to open the mouth and enter the gavage into the mouth and the esophagus and release the solution.

For mating, 5 female and 2 male mice were kept in one cage for 24 hours. Vaginal plaque was regarded as a sign of pregnancy. After mating, they were separated and tested in different weeks of pregnancy. One hundred and twenty pregnant mice were divided into four equal groups. The control group was without any intervention. Cesarean section was undertaken after one, two and three weeks of pregnancy to assess the embryos. Experimental group 1 received a daily dose of 3,500 mg/kg of coriander extract in gavage form. The experimental group 2 was fed by 5000 mg/kg of barberry extract per day via gavage. Experimental group 3 received a daily dose of 6500 mg/kg of barberry extract by gavage.

Data were analyzed using SPSS software (Version 20, Chicago, IL, USA). After assessing the normal distribution of data using Kolmogorov-Smirnov test, the comparison of quantitative data in different groups at similar times was done by One-way ANOVA and Kruskal-Wallis test. Comparison of quantitative data at different times in each group was performed by Paired T test, while qualitative data was done using Wilcoxon signed rank test. The statistical significant level was defined to be less than 0.05.

Results

Comparison of fetal height and weight in the first, second and third week of gestation in experimental and control groups is presented in Table 1 showing that fetal height in the first week of pregnancy in experimental groups of 1, 2 and 3 decreased significantly when compared to the control group. The reduction was significantly more in experimental group 3 when compared with experimental groups 1 and 2 ($P < 0.05$). For fetal weight in the first week of pregnancy, there was a significant decrease in experimental group 1, 2 and 3 in comparison to the control group and the decrease was more prominent in experimental groups 2 and 3 when compared with experimental group 1 ($P < 0.05$).

Table 1: Comparison of fetal height and weight in the 1st, 2nd and 3rd week of gestation between experimental and control groups

Variable	Control	Experimental 1	Experimental 2	Experimental 3
Fetal height (first week)	1.94±0.52 ^c	1.25±0.26 ^b	1.007±0.67 ^b	0.42±0.09 ^a
Fetal weight (first week)	0.916±0.45 ^b	0.67±0.56 ^b	0.26±0.129 ^a	0.085±0.03 ^a
Fetal height (second week)	0.93±0.24 ^b	0.87±0.288 ^b	0.866±0.533 ^b	0.296±0.2 ^a
Fetal weight (second week)	2.25±0.353 ^c	1.223±0.183 ^b	1.193±0.141 ^b	0.146±0.042 ^a
Fetal height (third week)	2.25±1.06 ^c	1.19±0.141 ^b	0.433±0.05 ^a	0.116±0.0288 ^a
Fetal weight (third week)	0.23±0.18 ^a	0.186±0.026 ^b	0.173±0.171 ^a	0.134±0.07 ^a

a, b, c: P<0.05

Discussion

During the recent decade, the consumption of traditional medicine has increased (15). They can affect several organs including the brain (17), digestive system (18) and reproductive system (19), so women's diet has an important role in safety of pregnancy (20). Herbal consumption during pregnancy can lead to congenital defects and abortions (21), therefore, researchers have investigated on the effect of herbal extracts during pregnancy with more caution (21-23). In this study, the effect of coriander extract on fetal height and weight was investigated showing that fetal height in the first week of pregnancy in experimental groups 1, 2 and 3 had a significant decrease compared to the control group. This decrease was significantly more in experimental group 3 compared to the experimental groups 1 and 2 (P<0.05). It was demonstrated that the use of *Matricaria chamomilla* in Balb/C mouse reduced the fetal height in a dose dependent manner (24) that was similar to our findings.

Identically, the use of ginger, which contains coriander-like materials including geraniol, flavonoids and tannins was shown to reduce the fetal height in a dose-dependent manner too. In fact, the geraniol in the coriander affects the dehydrogenase activity (25). Study of teratogenic properties of similar plants, such as chamomile, showed that flavonoids are responsible for these malformations. Also, chamomile compounds, similar to coriander, have the ability to attach to mucopolysaccharides of the cartilage tissue and change the appearance of the cartilage and the height of the fetus (25, 26) the same as our results.

Coriander has antioxidant activity and reduces the action of vitamin C (12). Considering that the

reduction of vitamin C activity has a prominent role on bone density and osteogenesis, coriander has a negative effect on fetal height. Similar findings were noted in our study. Evaluating the weight of embryos in the first, second and third week of pregnancy revealed that experimental groups showed a significant decrease compared to the control group (P<0.05). Tannin (11) and phenolic compounds in coriander were illustrated to increase the excretion of proteins, amino acids and essential amino acids (27). Tannin consumption sometimes leads to a severe poisoning and even death (28). Also, coriander extract was displayed to affect the metabolism of carbohydrates by increasing the activity of glycogen synthase, increasing liver glycogen concentration, inhibition of gluconeogenic and glycogen phosphorylase enzymes and decreases the level of glucose in the blood, which can reduce the weight of the fetus (29). Coriander extract with an anti-proliferative property and flavonoid contents can induce cell death and weight loss (30-32).

Conclusion

Coriander (*C. Sativum*) aqueous extract can cause a reduction in fetal height and weight in a dose-dependent manner during pregnancy. These findings can open a new window and concern for herbal medicines used during pregnancy.

Acknowledgement

The authors would like to thank Shiraz Branch, Islamic Azad University for financial support.

Conflict of Interest

None declared.

References

1 Mehrabani D, Zabeti M, Vahdati A, et al. The

- effect of aqueous extract of berberis vulgaris on fetal height and weight during pregnancy. *Int J Nutr Sci.* 2019;4:82-6. DOI:10.30476/IJNS.2019.82444.1020.
- 2 Tabei SMB, Tamadon A, Rahmanifar F, et al. Xenograft of mus musculus ovary in the uterus of rattus norvegicus after induction of pseudopregnancy. *Comp Clin Pathol.* 2016;25:1065-70. DOI:10.1007/s00580-016-2314-1.
 - 3 Hosseini SE, Mehrabani D, Avazpour Y, et al. Comparison of currant hydroalcoholic extract and diazepam on serum corticosterone levels in adult male mice. *Exp Zool J.* 2014;3:49-56.
 - 4 Rabbani Haghighi N, Mehrabani D. The protective effect of boiled coffee on liver enzymes AST, ALT, ALP and total bilirubin in rats treated with thioacetamide. *J Babol Univ Med Sci.* 2014;16:41-9.
 - 5 Hosseini SE, Khosrofarid M, Mehrabani D, et al. Perinatal and neonatal effects of rhizome extract of ginger on levels of insulin and ALT, AST, ALP on adult children of first-generation female rats. *J North Khorasan Univ.* 2015;7:292-307.
 - 6 Rezaei SMA, Nowrouzi M. Nutritional Patterns in Pregnant Women Referred to Yasuj Health Care Centers. *Int J Nutr Sci.* 2017;2:224-8.
 - 7 Jafarzadrh L, Asgari A, Golshan-Iranpoor F, et al. Abortifacient effects of stachys lavandulifolia vahl in mice. *J Shahrekord Univ Med Sci.* 2010;11:26-31.
 - 8 Zanganeh A, Rozbahani S, Monjemi R. Examining effect of the aqueous extract of anchusa italica on estrogen and progesterone concentrations and placental parameters in wistar female mice. *J Animal Physiol Dev.* 2016;9:49-60.
 - 9 Samavati R, Ducza E, Hajagos-Tóth J, et al. Herbal laxatives and antiemetics in pregnancy. *Reprod Toxicol.* 2017;72:153-8. DOI:10.1016/j.reprotox.2017.06.041. PMID:28610933.
 - 10 Naderi-Farsani M, Hoseinifar SH, Rashidian G, et al. Dietary effects of Coriandrum sativum extract on growth performance, physiological and innate immune responses and resistance of rainbow trout (Oncorhynchus mykiss) against Yersinia ruckeri. *Fish Shellfish Immunol.* 2019;91:233-40. DOI:10.1016/j.fsi.2019.05.031. PMID:31102711.
 - 11 Crespo YA, Bravo Sánchez LR, Quintana YG, et al. Evaluation of the synergistic effects of antioxidant activity on mixtures of the essential oil from Apium graveolens L., Thymus vulgaris L. and Coriandrum sativum L. using simplex-lattice design. *Heliyon.* 2019;5:e01942. DOI:10.1016/j.heliyon.2019.e01942. PMID:31245650.
 - 12 Akhlaghi M, Foshati S. Bioavailability and Metabolism of Flavonoids: A Review. *Int J Nutr Sci.* 2017;2:180-4.
 - 13 Akhlaghi M, Kohanmoo A. Mechanisms of Anti-Obesity Effects of Catechins: A Review. *Int J Nutr Sci.* 2018;3:127-32.
 - 14 Shoskes DA and et al. Quercetin in men with category III chronic Prostatitis: a preliminary prospective, double-blind, placebo-controlled trial. *Urology.* 1999;54:960-3. DOI:10.1016/s0090-4295(99)00358-1. PMID:10604689.
 - 15 Mehrabani G, Aminian S, Mehrabani G, et al. Dietetic plans within the multiple sclerosis community: A review. *Int J Nutr Sci.* 2019;4:14-22. DOI:10.30476/IJNS.2019.81531.1007.
 - 16 Dhiman S, Mukherjee G, Singh AK. Recent trends and advancements in microbial tannase-catalyzed biotransformation of tannins: a review. *Int Microbiol.* 2018;21:175-95. DOI:10.1007/s10123-018-0027-9. PMID:30810902.
 - 17 Aqababa H, Mottahedzadeh R, Shahabi S, et al. The action of ginkgo biloba on passive avoidance learning in experimental rats. *Middle East J Sci Res.* 2013;16:751-5.
 - 18 Rahnema M, Mehrabani D, Japoni S, et al. The healing effect of licorice (glycyrrhiza glabra) on helicobacter pylori infected peptic ulcers. *J Res Med Sci.* 2013;18:532-3. PMID:24250708.
 - 19 Zhu X, Proctor M, Bensoussan A, et al. Chinese herbal medicine for primary dysmenorrhoea. *Cochrane Database Syst Rev.* 2008;16:CD005288. DOI:10.1002/14651858.CD005288.pub3. PMID:18425916.
 - 20 Magee L, Vohra S, Matsui D, et al. Mother Nature: Establishing a Canadian Research Network for Natural Health Products (NHPs) During Pregnancy and Lactation. *J Altern Complement Med.* 2008;14:369-72. DOI:10.1089/acm.2007.0696. PMID:18466066.
 - 21 Marco DE, Elena C, Maria SL, et al. Prenatal drug exposure and teratological risk: One-year experience of an Italian teratology information service. *Med Sci Monit.* 2008;14:1-8. PMID:18227771.
 - 22 Dabirifard M, Maghsoudi Z, Dabirifard S, et al. Frequency, causes and how to use medicinal herbs during pregnancy. *Iran J Obstet Gynecol Infertil.* 2017;20:66-75.
 - 23 Parivar K, Kouchesfahani MH, Mohammadzadehasl B. Surveying Acyclovir side effects on developing embryos of Balb/C mice strain. *Med Sci J Islamic Azad Univ.* 2007;16:4:231-5.
 - 24 Baharara J, Rostampour M. Evaluation of teratogenic effects of Matricaria chamomilla

- in Balb/C mouse. *Iran J Obstet Infertil.* 2007;9:33-8.
- 25 Shariatzadeh MS, Azarnia M, Kaka GH, et al. Study of teratogenic effects of extract of berberis integerrima on liver tissue in mouse embryo. *J Cell Tissue.* 2013;4:63-70.
- 26 Sekiwa-Iijima Y, Aizawa Y, Kubota K. Geraniol dehydrogenase activity related to aroma formation in ginger (*Zingiber officinale* Roscoe). *J Agric Food Chem.* 2001;49:5902-6. DOI:10.1021/jf0102970. PMID:11743782.
- 27 Keating A, Chez RA. Ginger syrup as an antiemetic in early pregnancy. *Altern Ther. Altern Ther Health Med.* 2002;8:89-91. PMID:12233808.
- 28 Bravo L. Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance. *Nutr Rev.* 2002;56:317-3.
- 29 Deshpande SS. Handbook of food toxicology. Toxicants and antinutrient in plant foods. Marcel Dekker, New York; USA; 2002.p.920.
- 30 Chithra V, Leelamma S. Coriandrum sativum effect on lipid metabolism in 1,2-dimethyl hydrazine induced colon cancer. *J ethnopharmacol.* 2000;71:457-63. DOI:10.1016/S0378-8741(00)00182-3. PMID:10940583.
- 31 Nakano Y, Matsunaga H, Saita T, et al. Antiproliferative constituents in Umbelliferae Plants II. Screening for polyacetylenes in some Umbelliferae plants and isolation of panaxynol and falcarindiol from the root of *Heracleum moellendorffii*. *Biol Pharm Bull.* 1998;21:257-61. DOI:10.1248/bpb.21.257. PMID:9556156.
- 32 Blumenthal M, Goldberg A, Brinkmann J. Coriander seed. In: Herbal Medicine- Expanded Commission E Monographs, Integrative Medicine Communications, Newton, MA, USA; 2000.p.75-77.